5.1 Systematic Organic in Context

This first lesson topic takes an overview of the Organic reactions met in this and previous courses and some of the contexts in which these reactions are met.

Previous Chemistry

This activity examines the systematic approach to the reactions met in previous courses.



Learn:

- ***Names*** - ***RULES for naming !***
- ***Structures*** of both reactant molecule and product molecule
- ***Title*** of the reaction
- ***Reagents*** used to carry out the reaction

Substitution - an *atom* or *group* will be ***rem*** from a normally ***sat*** molecule to allow a ***diff*** *atom* or *group* to take its place.

Normally, the product formed is also ***sat*** and ***reac*** molecules react to form ***pro*** molecules

Equation using systematic names:

methane + $\longrightarrow$ *chloromethane* +

Equation using full structural formulae:

$$\begin{array}{ccc} & H & \\ & | & \\ H- & C & -H \\ & | & \\ & H & \end{array} \quad + \qquad \longrightarrow \qquad\qquad + $$

Equation using shortened structural formulae:

$$CH_{4(g)} \quad + \qquad \longrightarrow \qquad\qquad +$$

Cracking - a *larger* ***sat*** molecule is *broken apart* to produce *smaller* molecules, at least one of which will be ***uns*** .

Normally, a ***cat*** will be used. Sometimes only a couple of *neighbouring* ***hyd*** atoms will be '*cracked*' off to produce a single ***uns*** product. This reaction can also be called ***elimination***.

Equation using systematic names:

propane ⟶ *propene* +

Equation using full structural formulae:



⟶ +

Equation using shortened structural formulae:

$C_3H_{8(g)}$ ⟶ +

Dehydration - a specific ***elim*** reaction in which *neighbouring* ***hyd*** atom and ***hydroxyl*** group (—OH) will be '*cracked*' off to produce an ***uns*** product. The eliminated atoms form a stable ***wa*** molecule.

Equation using systematic names:

propan-1-ol ⟶ *propene* +

Equation using full structural formulae:



⟶ +

Equation using shortened structural formulae:

$C_3H_7OH_{(l)}$ ⟶ +

Elim is the reverse reaction to ***add*** .

Addition - a small molecule reacts with an ***uns*** molecule and adds *across* the ***dou*** bond to make a ***sat*** product.

Many different molecules can be added and many of these reactions have their own names.

Hydrogenation - add of *hyd*

Equation using systematic names:

propene + *hydrogen* $\xrightarrow{\text{Ni } catalyst}$

Equation using full structural formulae:



Equation using shortened structural formulae:

$C_3H_{6(g)}$ + $\longrightarrow$

Hydr is mainly used to convert *highly* ***unsat*** ***oils*** into *more* ***sat*** ***fats***. E.g. *vegetable oil* can be thickened and solidified to make *margarine* by ***hydro*** .

Halogenogenation - add of *hal*

Equation using systematic names:

propene + $\longrightarrow$ ***1,2-dibromopropane***

Equation using full structural formulae:



Equation using shortened structural formulae:

$C_3H_{6(g)}$ + $\longrightarrow$

The ***add*** reaction with a ***hal*** , usually ***bro*** , remains the accepted test for ***unsat*** - the presence of a C = C ***dou*** or C $\equiv$ C ***tri*** bond.

The ***hal*** is decol

bro , ***or*** $\longrightarrow$ *colourless*
chlo , ***gr*** $\longrightarrow$ *colourless*
io , ***br*** $\longrightarrow$ *colourless*

Hydrohalogenation - add of ***hyd*** ***hal***

Equation using systematic names:

propene + *hydrogen iodide* ⟶ ***1-iodo***

Equation using full structural formulae:



+ ⟶

Equation using shortened structural formulae:

$C_3H_{6(g)}$ + ⟶

Hydr is an alternative when only ***one halo*** atom is wanted on the product molecule. Depending on the position of the ***dou*** ***bond***, however, more than one ***iso*** is possible. In the above example, *2-iodopropane* is another possible product.



Hydration - add of ***wa***

Equation using systematic names:

propene + ⟶ ***propan-1-ol***

Equation using full structural formulae:



+ ⟶

Equation using shortened structural formulae:

$C_3H_{6(g)}$ + ⟶

Hydr of an ***alk*** is an important method for making *alcohols* but, like the previous example, more than one product can be formed.



In this case, the second ***iso*** would be *propan-2-ol*.

Propan-1-ol is a **primary alcohol** while *propan-2-ol* is a **secondary alcoho**l. The significance of these labels will become clearer in later lessons.

Condensation - two smaller molecule react to ***join together*** and form a larger molecule, ***eliminating*** a *small stable molecule*, usually ***water*** through one molecule losing ***an —H atom*** whilst the other molecule loses ***an —OH*** (*hydroxyl*) group.

Equation using systematic names:

propanoic acid + *ethanol* ⟶

Equation using full structural formulae:



Equation using shortened structural formulae:

$C_2H_5COOH_{(l)}$ + ⟶

The other main ***cond*** reaction that you will meet will involve an *acid* reacting with an *amine* to form an *amide*. This is very similar to the *ester* forming reaction above.



This is the reaction used to ***join*** many *amino acids* together to form *proteins*.

New Chemistry

This activity looks at some of the new reactions / families that you will meet in this Unit.



Much of the new Chemistry revolves round learning ***Oxi*** in more detail.

As can be seen, there are 3 types of ***Alc*** to be learnt and they have different products - ***Ald*** or ***Ket*** , or none - when oxidised.

Ald are the intermediates formed when a ***Prim*** ***Alc*** is oxidised to eventually form an ***Ac*** .

Cond becomes even more important - along with the reverse reaction, ***Hyd*** .

Not just ***Est*** are formed by a ***Cond*** reaction - ***Prot*** and ***Fa*** ***& Oi*** are also formed by ***Cond*** and broken down by the ***Hyd*** reaction.

So are the ***Sa*** of *Fatty Acids* formed when ***Alk*** ***Hyd*** is done to a ***Fat*** or ***Oil***.

Ter are natural *hydrocarbons*, found mainly in plants, formed by joining together smaller molecules called ***isoprene***, *2-methylbuta-1,3-diene*.

Terp belong to a group of chemicals usually referred to as *essential oils* which should not be mixed up with *edible oils* (***Fats & Oils***) and *mineral oils* derived from *Crude Oil* and made by the *Petrochemical* industry.

Functional Groups & Properties

This activity explains the functional groups found in these molecules and their possible effect on the chemical and physical properties of the molecule.

Family Name	*Functional Group*	*Name of Group*	*Chemical Reactions*	*Intermolecular Forces*	*Solubility in Water*
		carbon to carbon single		london dispersion forces	
		carbon to carbon double			
			oxidation condensation		very soluble
aldehyde					limited solubility
			none	polar-polar	
					
		carboxylate group			insoluble
amine					
		amide link		hydrogen bonding	only small amides soluble

As well as determining their ***Chem*** ***Reactions***, ***Func*** ***Groups*** can also effect the ***Phy*** ***Properties***. Properties such as



Acids Alcohols Amines	***Aldehydes Ketones Esters***	***Aromatics Alkenes Alkanes***

For example, the *vanilla bean* produces a compound called *vanillin*, which is used as a flavouring additive in sweet foods such as ice cream.



Vanillin

This molecule has effectively, 4 ***func*** ***groups***:-

The ***ben*** ***ring*** and the *ether* group (—O—CH_3) are dealt with in *Advanced Higher*.

The ***hydr*** group (—OH) and the ***carb*** group (—CH=O) will be expected to be learnt well this year.

Within the same molecule there can be ***non-po*** ***groups*** such as the ***aro*** ***ben*** ring, whilst the *ether* and ***ald*** (***carb*** are ***sli*** ***polar***.

Probably the most influential group will be the ***very po*** ***hydr*** group which is capable of ***hydr*** ***bonding*** and may make this molecule ***water sol*** .

Context - Kitchen Chemistry

This activity demonstrates how much of the Organic Chemistry met in this Unit will be taught within the context of Kitchen Chemistry