


Higher Chemistry



Topic 4:

Enthalpy, Moles & RedOx

Answer Book

4.1	A	B	C	D
1				
2				
3				
4				
5				
6				
7				

4.2 4.3	A	B	C	D
1				
2				
3				
4				
5				
6				
7				
8				

4.4	A	B	C	D
1				
2				
3				
4				
5				
6				
7				

4.5 4.6	A	B	C	D
1				
2				
3				
4				
5				
6				
7				
8				

4.7	A	B	C	D
1				
2				
3				
4				
5				
6				
7				

4.8	A	B	C	D
1				
2				
3				
4				
5				
6				
7				
8				

4.9 4.10	A	B	C	D
1				
2				
3				
4				
5				
6				

HOME PRACTICE ANSWERS

4.1

- Q1.** The standard enthalpy of combustion of propane is $-2219 \text{ kJ mol}^{-1}$.
- a)** C_3H_8 . 1
- b)** $\text{C}_3\text{H}_{8(g)} + 5 \text{O}_{2(g)} \rightarrow 3 \text{CO}_{2(g)} + 4 \text{H}_2\text{O}_{(l)}$ 1
- c)** Molar mass of $\text{C}_3\text{H}_8 = (3 \times 12) + (8 \times 1) = 44\text{g}$ (1)
- $-2219 / 44 = -50.4 \text{ kJ g}^{-1}$ (1) 2
- d)** $-50.4 \text{ kJ} \times 1000 = 50,400 \text{ kJ kg}^{-1}$ 1
- Q2. a)** any *two* from:
- (supports hypothesis) because when the fuel contains more carbon the temperature of the water went up more / faster (in 2 minutes)
- (does not support hypothesis as) temperature change per gram decreases as the number of carbons increases
- (does not support hypothesis) because the more carbon in the fuel the more smoke or the dirtier / sootier it is 2
- b)** The student only tested hydrocarbons / alkanes / fuels with between 5 and 12 carbon atoms so
- should have tested molecules from other families 1
- Q3.** Large amount of energy released / very exothermic (1)
- High proportion (all) of products are gases - propulsion (1) 2
- Total** (10)

HOME PRACTICE ANSWERS

4.2/4.3

Q1. $\Delta H = (\text{Energy of Bond Breaking}) + (\text{Energy of Bond Forming})$ (½)

Bond Breaking = (4 x $\Delta H(\text{C—H}) + \Delta H(\text{C=C}) + \Delta H(\text{H—Cl})$) (½)
 (endothermic) = (4 x 467) + (612) + (432) = + 2912 kJ (½)

Bond Forming = (5 x $\Delta H(\text{C—H}) + \Delta H(\text{C—C}) + \Delta H(\text{C—Cl})$) (½)
 (exothermic) = (5 x 467) + (347) + (346) = - 3028 (½)

overall ΔH = (+2912) + (-3028)
 = - 116 kJ mol⁻¹ (½)

3

Q2. a) no effect

1

b) $E = m c \Delta T$
 = 0.1 (½) x 4.18 x 16 = 6.69 kJ (½)

1000 cm³ of H₂O_{2(aq)} ↔ 0.88 mol

50 cm³ of H₂O_{2(aq)} ↔ 50/1000 x 0.88 (½)
 = 0.044 mol (½)

0.044 mol ↔ 6.69 kJ

1 mol ↔ 1/044 x 6.69 = 152 kJ (½)

include negative sign in final answer -152 kJ (½)

(units not required; deduct ½ mark for incorrect units)

3

Q3. a) $E = m c \Delta T$
 = 2 x 4.18 x 75 (½) = 627 kJ (½)

Data Book, Butane 2878 kJ ↔ 1mol (½)
 627 kJ ↔ 1 x 627 / 2878 = 0.218 mol (½) 2

b) 80% ↔ 0.218 mol (½)
 100% ↔ 100/80 x 0.218 = 0.2725 mol (½)


Butane, C₄H₁₀, 1 mol ↔ 58g (½)

0.2725 mol ↔ 0.2725/1 x 58 = 15.8 g (½) 2


Total (11)

HOME PRACTICE ANSWERS


4.4




(3 'sensible' numbers required for $\frac{1}{2}$ mark for addition based on following through;
 no units required; deduct $\frac{1}{2}$ for incorrect units)



(deduct $\frac{1}{2}$ mark for incorrect addition based on numbers used; no units required;
 deduct $\frac{1}{2}$ mark for incorrect units)



(3 'sensible' numbers required for $\frac{1}{2}$ mark for addition based on following through;
 no units required; deduct $\frac{1}{2}$ mark for incorrect units)







(3 'sensible' numbers required for $\frac{1}{2}$ mark for addition based on following through;
 no units required; accept kJ; deduct $\frac{1}{2}$ mark for incorrect units)




Total (8)

HOME PRACTICE ANSWERS

4.5/4.6



(deduct ½ mark for no or incorrect units)










(deduct half mark for missing or incorrect unit) **2**

Total (10)

HOME PRACTICE ANSWERS

4.7



so Cl_2 is limiting reagent . **3**




Total **(10)**

HOME PRACTICE ANSWERS

4.8

Q1. a)	i)	$\text{Mg}_{(s)} \rightarrow \text{Mg}^{2+}_{(aq)} + 2e^{-}$	(1/2)	
	ii)	$\text{Sn}^{4+}_{(aq)} + 2e^{-} \rightarrow \text{Sn}^{2+}_{(aq)}$	(1/2)	
	iii)	$\text{ClO}^{-}_{(aq)} + 2\text{H}^{+}_{(aq)} + 2e^{-} \rightarrow \text{Cl}^{-}_{(aq)} + \text{H}_2\text{O}_{(l)}$	(1/2)	
	iv)	$\text{FeO}^{2-}_{4(aq)} + 8\text{H}^{+}_{(aq)} + 3e^{-} \rightarrow \text{Fe}^{3+}_{(aq)} + 4\text{H}_2\text{O}_{(l)}$	(1/2)	2
b)	i)	Oxidation	(1/2)	
	ii)	Reduction	(1/2)	
	iii)	Reduction	(1/2)	
	iv)	Reduction	(1/2)	2
Q2. a)	i)	$\text{Zn}_{(s)} \rightarrow \text{Zn}^{2+}_{(aq)} + 2e^{-}$	(1/2)	
	ii)	$2\text{Al}_{(s)} \rightarrow 2\text{Al}^{3+}_{(aq)} + 6e^{-}$	(1/2)	
	iii)	$2\text{I}^{-}_{(aq)} \rightarrow \text{I}_{2(s)} + 2e^{-}$	(1/2)	
	iv)	$\text{Zn}_{(s)} \rightarrow \text{Zn}^{2+}_{(aq)} + 2e^{-}$	(1/2)	2
b)	i)	$\text{Pb}^{2+}_{(aq)} + 2e^{-} \rightarrow \text{Pb}_{(s)}$	(1/2)	
	ii)	$6\text{H}^{+}_{(aq)} + 6e^{-} \rightarrow 3\text{H}_{2(g)}$	(1/2)	
	iii)	$\text{Cl}_{2(g)} + 2e^{-} \rightarrow 2\text{Cl}^{-}_{(aq)}$	(1/2)	
	iv)	$\text{S}_{(s)} + 2e^{-} \rightarrow \text{S}^{2-}_{(s)}$	(1/2)	2
Q3.		$\text{Cd}_{(s)} + 2\text{OH}^{-}_{(aq)} \rightarrow \text{Cd}(\text{OH})_{2(s)} + 2e^{-}$		
		$\text{Ni}(\text{OH})_{3(s)} + e^{-} \rightarrow \text{Ni}(\text{OH})_{2(s)} + \text{OH}^{-}_{(aq)}$		<i>multiply by 2</i>
a)	$\text{Cd}_{(s)} + 2\text{Ni}(\text{OH})_{3(s)} \rightarrow \text{Cd}(\text{OH})_{2(s)} + 2\text{Ni}(\text{OH})_{2(s)}$			1
b)	$\text{Ni}(\text{OH})_{3(s)}$			1
				Total (10)

HOME PRACTICE ANSWERS**4.9/4.10**

(State symbols not required; accept $\text{I}_{2(\text{s})}$ on right hand side of equation; deduct ½ if 2e^- shown on each side).

b) no. of moles of thiosulphate = $0.0050 \times 0.0149 = 7.45 \times 10^{-5}$ (½)

2 mol thiosulphate : 1 mol I_2 (1)

no. of moles of $\text{I}_2 = \frac{1}{2} \times 7.45 \times 10^{-5} = 3.725 \times 10^{-5}$ (½)

1 mol $\text{I}_2 = 2 \times 126.9 = 253.8$ g (½)

mass of $\text{I}_2 = 3.725 \times 10^{-5} \times 253.8 = 0.00945$ g (9.45×10^{-3} g) (½)

(Deduct ½ for no or incorrect units) **3**



(state symbols not required)

b) there is a colour change from colourless to purple (or purple to colourless)

or the reaction is self-indicating (or a colour change shows the end of the reaction) **1**

c) i) first titre is a rough (or approximate) result
 or not accurate
 or an estimate
 or too far away from the others **1**

ii) no. of moles of $\text{MnO}_4^-_{(\text{aq})} = 0.040 \times 0.0269 = 0.001$ (½)

ratio of $(\text{COOH})_{2(\text{aq})} : \text{MnO}_4^-_{(\text{aq})} = 5 : 2$ (½)

no. of moles of $(\text{COOH})_{2(\text{aq})}$ in 25 cm³ = $5/2 \times 0.001 = 0.0025$ (½)

no. of moles of $(\text{COOH})_{2(\text{aq})}$ in 500 cm³ = $0.0025 \times 500/25 = 0.05$ (½)

(no units required; deduct ½ mark for incorrect units) **2**

Total (9)

CONSOLIDATION ANSWERS

A

- Q1.** a) 1130 kJ 1
 b) - 640 kJ 1
 c) Exothermic (½)
 Because ΔH is negative.
 or Products have less energy than reactants (½) 1
- Q2.** a) Loss of heat (½) to beaker/ thermometer/ air/ surroundings (½) 1
 b) Any **one** from:
 Use plastic / better insulated beaker.
 Use a vacuum flask for reaction.
 Have a lid on top (to reduce heat loss).
 Use a more accurate thermometer. 1
- Q3.** NH_3 , 1 mole = 17g $3.4\text{g} = 3.4/17 = 0.2 \text{ mol of NH}_3$ (½)
 1 mol NH_3 \rightarrow 3 mol of N atoms
 0.2 mol of NH_3 \rightarrow 0.6 mol of N atoms C
 1 mol \rightarrow 6.02×10^{23} N atoms (½)
 0.6 mol \rightarrow $0.6 \times 6.02 \times 10^{23} = 3.612 \times 10^{23}$ N atoms (½) 2
- Q4.** a) $\text{C}_3\text{H}_{8(g)}$ \rightarrow 3 $\text{CO}_{2(g)}$
 1 mol \rightarrow 3 mol (½)
 0.2 mol \rightarrow **0.6 mol** (½) 1
- b) 1 mol \rightarrow 23 l (½)
 0.6 mol \rightarrow $0.6 \times 23 = 13.8 \text{ l}$ (½) 1
- Q5.** a) $\text{N}_{2(g)}$ + 2 $\text{O}_{2(g)}$ \rightarrow 2 $\text{NO}_{2(g)}$
 1 mol \quad 2 mol
 1 vol \quad 2 vol (½)
- So 80 l of N_2 would require 160 l of O_2 (½)- only 20 l available so O_2 is limiting reagent
 20 l of O_2 would react with 10 l of N_2 (½) so $80 - 10 = 70 \text{ l of N}_2$ in excess. (½) 2
- b) $\text{N}_{2(g)}$ + 2 $\text{O}_{2(g)}$ \rightarrow 2 $\text{NO}_{2(g)}$
 \quad 2 mol \quad 2 mol
 \quad 2 vol \quad 2 vol (½)
 \quad 20 l \quad 20 l (½)
- plus 70 l of unreacted N_2 (½), so total volume = $70 + 20 = 90 \text{ l of gas}$. (½) 2

CONSOLIDATION ANSWERS**A**


- Q6. *a)* $I_{2(aq)} + 2e^- \rightarrow 2I^-_{(aq)}$ **1**
- b)* $C_6H_8O_{6(aq)} + I_{2(aq)} \rightarrow 2H^+I^-_{(aq)} + C_6H_6O_{6(aq)}$
- or*
- $C_6H_8O_{6(aq)} + I_{2(aq)} \rightarrow 2H^+_{(aq)} + 2I^-_{(aq)} + C_6H_6O_{6(aq)}$ **1**
- c)* Reducing agent **1**
- Q7. *a)* None *or* no effect **1**
- b)* Any purple colour would disappear. **1**

CONSOLIDATION ANSWERS

B


- Q1. a)** The overall enthalpy change in a reaction (or sequence of reactions) depends only on the reactants and the products and not on the route taken. **1**
- b)** Combustion of magnesium sulphide
- or* $\text{MgS} + \frac{3}{2}\text{O}_2 \rightarrow \text{MgO} + \text{SO}_2$
- or* $2\text{MgS} + 3\text{O}_2 \rightarrow 2\text{MgO} + 2\text{SO}_2$ **1**
- b)** (ΔH for the) combustion of magnesium sulphide. **1**

Q2. Ammonium sulphate fertiliser can be manufactured by neutralising sulphuric acid with ammonia.





- a)** 132 g \rightarrow 1 mol $(\text{NH}_4)_2\text{SO}_{4(s)}$ \rightarrow 2 mol NH_4^+ ions ($\frac{1}{2}$)
- 1 mol \rightarrow 6.02×10^{23} ions 2 mol \rightarrow 1.204×10^{24} ions ($\frac{1}{2}$) **1**

b) *though 20 tonnes (20,000,000 g) and 49 tonnes (49,000,000 g) could be converted into grammes and actual numbers of moles calculated, it is the ratio that counts and both calculations can be done with original numbers retained.*



Since H_2SO_4 has the lower mole ratio it is the limiting reagent and will run out first. **2**

An alternative method is to work out the amount of product produced by each chemical and whichever produces the least is the limiting reagent.



CONSOLIDATION ANSWERS

B

- Q3.** $\text{H}_{2(g)} + \frac{1}{2}\text{O}_{2(g)} \rightarrow \text{H}_2\text{O}_{(l)}$ (state symbols *are* required) **1**
- Not* $2\text{H}_{2(g)} + \text{O}_{2(g)} \rightarrow 2\text{H}_2\text{O}_{(l)}$
- Q4. a)**
- Initial temperature of water (½)
 - Final (or maximum) temperature of water (½)
 - Initial mass of ethanol (+ container) (½)
 - Final mass of ethanol (+ container) (½) **2**
- b) i)** -1367 kJ mol⁻¹ (-½ if unit incorrect or missing) **1**
- ii)** Heat loss to air **or** Heat loss to beaker **1**
- [*not* inaccuracy in thermometer or balance]
- Q5.**
- | | | | | | | | |
|-------------------------|-----|---------------|---------------|-------------------------|-----|--------------------------|--------------|
| Fe_2O_3 | $+$ | 3H_2 | \rightarrow | 2Fe | $+$ | $3\text{H}_2\text{O}$ | (½) |
| 1 mol | | 3 mol | | | | | (½) |
| | | | | 1 mol of H ₂ | = | 23.2 l | (½) |
| | | | | 3 mol of H ₂ | = | 3 x 23.2 = 69.6 l | (½) 2 |
- Q6. a)** The purple colour just remains / appears. **1**
- b)** $(15.8 + 15.6) / 2 = 15.7 \text{ cm}^3$ **1**
- Q7. a)**
- Dissolve the tablet in a little deionised water (½)
 - and make up exactly to the mark in a graduated flask. (½) **1**
- b)** Add a little starch solution to the flask. (½)
- The end point is when the mixture just turns a permanent blue/black colour. (½) **1**
- Q8. a)** $\text{ClO}_4^{-}{}_{(aq)} + 8\text{H}^{+}{}_{(aq)} + 8\text{e}^{-} \rightarrow \text{Cl}^{-}{}_{(aq)} + 4\text{H}_2\text{O}_{(l)}$ **1**
- b)** H⁺_(aq) ions are involved/ used up. **1**



CONSOLIDATION ANSWERS

C

Q1. a) $\Delta H = -cm \Delta T$ (1/2)
 $= -4.18 \times 0.2 \times (-2)$ (1/2)
 $= +1.7 \text{ kJ}$ (1) [-1/2 if no, or incorrect unit] **2**


b) Any **one** from:

no lid/cover to flask (to prevent convection)
 not waiting to ensure minimum temperature reached
 not ensuring that the vacuum flask was also at 24 °C **1**



Q3. Twice (as many methane molecules) **1**

Q4. Magnesium reacts with 1 mol l^{-1} hydrochloric acid to form hydrogen.




Q5. a) Endothermic. **1**




b) $\Delta H = -cm \Delta T$ (1/2)
 $= -4.18 \times 0.5 \times (-8)$ (1/2)
 $= +16.72 \text{ kJ}$ (1) [-1/2 if no, or incorrect unit] **2**

CONSOLIDATION ANSWERS


C




An alternative method is to work out the amount of product produced by each chemical and whichever produces the least is the limiting reagent.



Q8. An electrochemical cell in which the following changes took place was set up.



- a) The second equation must be tripled / multiplied by 3.
or Both of the equations must have an equal number of electrons. 1
- b) $\text{Cr}_2\text{O}_7^{2-}{}_{(aq)} + 14\text{H}^+{}_{(aq)} + 6\text{Br}^-{}_{(aq)} \rightarrow 2\text{Cr}^{3+}{}_{(aq)} + 7\text{H}_2\text{O}_{(l)} + 3\text{Br}_{2(aq)}$ 1
- c) $\text{Br}^- / \text{Br}^-{}_{(aq)} / \text{bromide (ion)} / \text{bromide solution}$ 1




- b) None / no effect
or Reduce the margin of error 1 increase the accuracy. 1


CONSOLIDATION ANSWERS

D


Q1. a)




$$\Delta H = - 882 \text{ kJ}$$



$$\Delta H = + 283 \text{ kJ}$$




$$\Delta H = + 858 \text{ kJ}$$




[Correct equations (1),
(1/2) if only one error]

[Correct ΔH values (1),
(1/2) if only one error]



So 80 cm³ of CO would require 40 cm³ of O₂ (1/2) - 150 cm³ available so O₂ is in excess and CO is the **limiting reagent**. (1/2)



80 cm³ of CO would **produce 80 cm³ of CO₂** (1/2) 3

Q3. a) 1 l of water = 1 kg (1/2)

$$\Delta H = - cm \Delta T \quad (1/2)$$

$$= -4.18 \times 1 \times (-2.5) \quad (1/2)$$

$$= +10.5 \text{ kJ} \quad (1/2) \quad [-1/2 \text{ if no, or incorrect unit}] \quad 2$$

b) KBr, 1 mol = 119 g (1/2)

$$238 \text{ g} \longrightarrow +10.5 \text{ kJ} \quad (1/2)$$

$$119 \text{ g} \longrightarrow +10.5 \times 119/238 \quad (1/2) = +5.25 \text{ kJ mol}^{-1} \quad (1/2) \quad 2$$

Q4. a) L (Data Booklet) 1

b) Al(NO₃)₃ so 3 mol of nitrate ions per mol of aluminium nitrate 1

c) MgSO₄ MgBr₂ (1/2)

4 mol of SO₄ means 4 mol of MgSO₄ \longrightarrow 4 mol Mg (1/2)

Therefore, Mg in MgBr₂ = 7 - 4 = 3 mol (1/2)

So Br in MgBr₂ = 3 x 2 = 6 mol (1/2) 2

CONSOLIDATION ANSWERS

D

- Q5. a)** It is reduced / receives electrons / gains electrons. **1**
- b)** $2 \text{Al}_{(s)} + 6 \text{H}^+_{(aq)} \rightarrow 2 \text{Al}^{3+}_{(aq)} + 3 \text{H}_{2(g)}$ **1**
- Q6. a)** Either of the following:
- The error in reading the larger volume is a smaller portion of the total volume.
- The error in reading the longer time is a smaller portion of the total time. **1**
- b)** The volume occupied by a gas changes with temperature. **1**
- Q7. a)** The addition of ions ($\frac{1}{2}$) from the potassium iodide ($\frac{1}{2}$) **1**
- b)** All of the vitamin C has been oxidised / reacted ($\frac{1}{2}$) so no more H^+ or I^- ions are being produced. ($\frac{1}{2}$) **1**